

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**

- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 467 349 A1**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **91111983.2**

(51) Int. Cl.<sup>5</sup>: **C12N 1/21, //C12N15/63**

(22) Date of filing: **19.10.84**

This application was filed on 17 - 07 - 1991 as a divisional application to the application mentioned under INID code 60.

(30) Priority: **20.10.83 US 543528**  
**01.03.84 US 585282**

(43) Date of publication of application:  
**22.01.92 Bulletin 92/04**

(60) Publication number of the earlier application in accordance with Art.76 EPC: **0 140 308**

(84) Designated Contracting States:  
**AT BE CH DE FR GB IT LI LU NL SE**

(71) Applicant: **THE RESEARCH FOUNDATION OF STATE UNIVERSITY OF NEW YORK**  
**State University Plaza, Broadway**  
**Albany, New York 12246(US)**

(72) Inventor: **Inouye, Masayori**  
**24 Hilltop Lane**  
**Setauket New York 11733(US)**  
Inventor: **Mizuno, Takeshi**  
**9-105 Popuragoaka 2-11-3 Naruse**  
**Machida-shi Tokyo 194(JP)**  
Inventor: **Chou, Mel-Yin**  
**668 Hawkins Road East**  
**Coram New York 11727(US)**

(74) Representative: **Vossius & Partner**  
**Siebertstrasse 4 P.O. Box 86 07 67**  
**W-8000 München 86(DE)**

(54) Regulation of gene expression by employing translational inhibition utilizing mRNA interfering complementary RNA.

(57) Disclosed are organisms containing a nucleic acid construct, said nucleic acid construct being an artificial nucleic acid construct which, upon introduction into a cell of said organism containing a gene, regulates the function of said gene and produces a ribonucleotide sequence which does not naturally occur in the cell, is complementary to at least a portion of a ribonucleotide sequence transcribed by said gene, and regulates the function of said gene.

**EP 0 467 349 A1**

The control or regulation of the gene expression of the genetic material of cellular material or an organism has received special attention by scientists and in special circumstances, employing recombinant DNA and other techniques, has been achieved. For example, in the PCT Patent Application WO 83/01451 published April 23, 1983, there is disclosed a technique employing an oligonucleotide, preferably in phosphotriester form, having a base sequence substantially complementary to a portion of messenger ribonucleic mRNA coding for a biological component of an organism. This oligonucleotide is introduced into the organism and, due to the complementary nature of the oligonucleotide and the messenger ribonucleotide, the two components hybridize under appropriate conditions to control or inhibit synthesis of the organism's biological component coded for by the messenger ribonucleotide. If the biological component is vital to the organism's viability, then the oligonucleotide could act as an antibiotic. A related technique for the regulation of gene expression in an organism is described in an article appearing in *Cell*, Vol. 34, p. 683 of September, 1983. The disclosures of the above-identified publications are herein incorporated and made part of this disclosure.

As indicated hereinabove, it is known that the expression of certain genes is capable of being regulated at the level of transcription. Transcriptional regulation is carried out either negatively (repressors) or positively (activators) by a protein factor. It is also known that certain specific protein factors regulate translation of specific mRNAs. Also, as indicated hereinabove, it has become evident that RNAs are involved in regulating the expression of specific genes and it has been reported that a small RNA transcript of 174 bases is produced, upon growing *Escherichia coli* in a medium of high osmolarity, which inhibits the expression of the gene for an outer membrane protein (OmpF) protein, see "Regulation of Gene Expression by a Small RNA Transcript (micRNA) in *E. coli* K12", *Proc. Jap. Acad.*, 59, 335-338 (1983). The inhibition of OmpF protein production by the small RNA transcript (micRNA, i.e. mRNA interfering complementary RNA) is likely due to the formation of the hybrid between the micRNA and the ompF mRNA over a region of approximately 80 bases including the Shine-Dalgarno sequence and the initiation codon. A similar regulation by a small complementary RNA has also been described for the Tn10 transposase, see Simons et al "Translational Control of IS10 Transposition", *Cell*, 34, 683-691 (1983). In this case, however, the gene for the transposase and the gene for the micRNA are transcribed in opposite directions off the same segment of DNA such that the 5'-ends of the transcripts can form a complementary hybrid. The hybrid is thought to inhibit translation of the transposase mRNA. However, the transposase situation is in contrast to the ompF situation in which the ompF gene and the micRNA gene (micF) are completely unlinked and map at 21 and 47 minutes, respectively, on the *E. coli* chromosomes.

It is an object of this invention to provide transformed organisms having special properties with respect to the gene expression of the genetic material making up said organisms.

How this object of the present invention is achieved will become apparent in the light of the accompanying disclosure and with reference to the accompanying drawings wherein:

Fig. 1 describes the construction of a subclone or a gene and various plasmids carrying the promoter region therefor;

Fig. 2 sets forth the nucleotide sequence of the promoter region and upstream of a gene, specifically the ompC gene;

Fig. 3 illustrates the hybrid formation between certain RNA in accordance with the practices of this invention;

Fig. 4 illustrates the homologous sequences between certain genes, specifically micF and the ompC genes; and

Fig. 5 illustrates a possible model for the role of RNA, specifically micF RNA useful in and in accordance with the practices of this invention.

Fig. 6 illustrates the construction of mic vector pJDC402 and mic(lpp).

Fig. 7 illustrates the homology between the ompC mRNA and the lpp mRNA; and wherein

Fig. 8 illustrates fragments used to construct mic(ompA) genes.

Gene expression of the genetic material of cellular material or an organism in accordance with the practices of this invention is regulated, inhibited and/or controlled by incorporating in or along with the genetic material of the cellular material or organism DNA or other genetic material which transcribes to an RNA which is complementary to and capable of binding or hybridizing to the mRNA of the genetic material of said organism or cellular material. Upon binding to or hybridization with the mRNA, the translation of the mRNA is prevented with the result that the product, such as protein material coded for by the mRNA is not produced. In the instance where the mRNA translated product, e.g. protein, is vital to the growth of the organism or cellular material, the organism or cellular material so transformed or altered becomes, at least, disabled.

In accordance with the practices of this invention there has been constructed a mic system designed to

regulate the expression of a gene. More particularly, one can construct in accordance with the practices of this invention an artificial mic system to regulate the expression of any specific gene in *E. coli*.

Further, in accordance with the practices of this invention, a micRNA system for a gene is constructed by inserting a small DNA fragment from the gene, in the opposite orientation, after a promoter. Such a system provides a way, heretofore unknown, for specifically regulating the expression of any gene. More particularly, by inserting the micDNA fragments under the control of an inducible promoter, particularly as embodied in *E. coli*, the expression of essential *E. coli* genes can be regulated. It would appear, therefore, that in accordance with the practices of this invention, the inducible lethality thus-created may be an effective tool in the study of essential genes.

Hereinafter, in accordance with the practices of this invention, there is described the construction of an artificial mic system and the demonstration of its function utilizing several *E. coli* genes. The mic system in accordance with this invention is an effective way to regulate the expression of specific prokaryotic genes. This invention accordingly provides the basis for accomplishing similar regulation of biologically important genes in eukaryotes. For example, the mic system can be used to block the expression of harmful genes, such as oncogenes and viral genes, and influence the expression of substantially any other gene, harmful or otherwise.

The practices of this invention are applicable to both procaryotic and eucaryotic cellular materials or micro-organisms, including bacteria, yeast and viruses, and is generally applicable to organisms, which contain genetic material which is expressed.

Accordingly, in the practices of this invention from a genetic point of view as evidenced by gene expression, new organisms are readily produced. Further, the practices of this invention provide a powerful tool or technique for altering gene expression of the genetic material making up organisms and the like so as to make such organisms disabled or incapable of functioning normally or imparting special properties thereto. The DNA material employed in the practices of this invention can be incorporated into the organisms to be treated or effected, such as by direct introduction into the nucleus of a eucaryotic organism or by way of a plasmid or suitable vector containing the special DNA of this invention in case of a procaryotic organism.

By way of further background of the practices of this invention, it has been found that the expression of the genes for the major outer membrane proteins, OmpF and OmpC, of *Escherichia coli* are osmoregulated. The *ompC* locus was found to be transcribed bidirectionally under conditions of high osmolarity, and the upstream transcript RNA of approximately 170 bases was found to inhibit the production of OmpF protein. This RNA (micRNA) has a long sequence which is complementary to the 5'-end region of the *ompF* mRNA that includes the ribosome-binding site and the coding region of the first nine amino acid residues of pro-OmpF protein. Thus, it is proposed that micRNA inhibits the translation of *ompF* mRNA by hybridizing with it. This novel mechanism can account for the observation that the total amount of the OmpF and of the OmpC proteins is always constant in *E. coli*.

The major outer membrane proteins of *Escherichia coli*, OmpF and OmpC, are essential proteins which function as passive diffusion pores for small, hydrophilic molecules. These matrix porin proteins are encoded by the structural genes *ompF* and *ompC*, which map at 21 and 47 min on the *E. coli* chromosome, respectively, see Reeves, P. in *Bacterial Outer Membranes: Biogenesis and Function* (ed. Inouye, M.) 255-291 (John Wiley and Sons, New York, 1979). The expression of these genes is regulated by the osmolarity of the culture medium. There is a strict compensatory production of both proteins: as the osmolarity of the culture medium increases, the production of OmpF protein decreases, while the production of OmpC protein increases so that the total amount of the OmpF plus OmpC proteins is constant. This osmoregulation of the *ompF* and *ompC* genes is controlled by another unlinked locus, *ompB*, which maps at 74 min, see Hall, M.N. & Silhavy, T.J., *J. Mol. Biol.* 146, 23-43 (1981) and Hall, M.N. & Silhavy, T.J., *J. Mol. Biol.* 151, 1-15 (1981). The *ompB* locus contains two genes called *ompR* and *envZ*. The DNA sequences of both genes have been determined and their gene products have been characterized, see Wurtzel, E.T. et al., *J. Biol. Chem.* 257, 13685-13691 (1982) and Mizuno, T., et al., *J. Biol. Chem.* 257, 13692-13698 (1982). The *EnvZ* protein is assumed to be a membrane receptor protein which serves as an osmosensor and transmits the signal from the culture medium to the OmpR protein. The OmpR protein then serves as a positive regulator for the expression of the *ompF* and *ompC* genes. The *ompF* and *ompC* genes were sequenced, and extensive homology was found in their coding regions, while there was very little homology in their promoter regions. It was during the course of the characterization of the *ompC* gene, that the novel regulatory mechanism of gene expression mediated by a new species of RNA called mRNA interfering complementary RNA (micRNA) in accordance with this invention was discovered and/or elicited. MicRNA is produced from an independent transcriptional unit (the *micF* gene). This gene is located immediately upstream of the *ompC* gene but is transcribed in the opposite direction. The 174-base micRNA blocks the

translation of the *ompF* mRNA by hybridizing to it. Since the production of micRNA is assumed to be proportional to the production of *ompC* mRNA, this regulatory mechanism appears to be a very efficient way to maintain a constant total amount of OmpF and OmpC proteins.

## A DNA Fragment Suppressing *ompF* Expression

While characterizing the *ompC* promoter, it was found that a DNA fragment of approximately 300 bp, located upstream of the *ompC* promoter, completely blocked the production of OmpF protein when OmpF<sup>+</sup> cells were transformed with a multi-copy plasmid harboring this DNA fragment. For this experiment, plasmid pMY150 was constructed from the original *ompC* clone, pMYIII, see Mizuno, T. et al, J.Biol. Chem. 258, 6932-6940 (1982), by changing the HpaI sites of pMYIII to HbaI sites followed by removal of the 1.1 kb Sall fragment as described in Fig. 1a of Fig. 1.

In Fig. 1 there is shown the construction of the subclone of the *ompC* gene and various plasmids carrying the *ompC* promoter region.

(a) Schematic presentation of the subcloning of the *ompC* gene. Plasmid pMYIII carrying a 2.7 Kb E. coli chromosomal DNA in pBR322 was described previously. The plasmid (1 ug of DNA) was digested with HpaI and religated in the presence of an XbaI linker (CTCTAGAG, 150 p mole). Thus, ca. 400 bp HpaI fragment was removed and a unique XbaI site was newly created (pMYI00). Plasmid pMYI00 (1 ug of DNA) was further digested with Sall and religated to remove a 1.1 kb Sall fragment (pMYI50). In order to obtain an *ompC* promoter fragment of different sizes, plasmid pMYI50 was digested by Bal 31 nuclease after cleavage of the unique BglII site (see Fig 1b), subsequently the plasmid was religated in the presence of an XbaI linker. Plasmid pCX28 thus constructed is one of clones carrying approximately 300-bp XbaI-XbaI fragment as shown in Fig. 1b.

(b) Simplified restriction map of the plasmid pMYI50 carrying the entire *ompC* gene. The 1.8 Kb HindIII-Sall fragment (boxed region) in pBR322 contains the entire *ompC* coding region as well as the 5'- and 3'-non-coding region. Transcription of the *ompC* gene proceeds in the direction shown by an arrow. A bidirectional arrow indicates an approximate deleted region (ca. 600 bp) for plasmid pCX28.

(c) various  $\beta$ -galactosidase (*lacZ*) gene fusions to the DNA fragments derived from the *ompC* promoter and its upstream region: Plasmid I, 507-bp XbaI-RsaI fragment was isolated from pMYI50 (an RsaI site is present just downstream of the ATG codon), and inserted between XbaI-SmaI sites of plasmid pICIII which is derived from plasmid pINIII carrying the *lacZ* gene. During the ligation, a HindIII linker was inserted between the RsaI and SmaI ligation site. The XbaI-HindIII fragment was isolated from the plasmid thus constructed and reinserted into plasmid pKM005 to create a *lacZ* gene fusion in the right reading frame. Characteristic features of plasmids pICIII and pKM005 were described previously. Plasmids II and IV carrying approximately 430-bp MspI-BamHI fragment was isolated from clone I (a BamHI site is present just downstream of the ATG codon for the  $\beta$ -galactosidase coding sequence in plasmid I), and treated with SI nuclease to create blunt ends. After adding XbaI linkers at both ends, the XbaI-XbaI fragment thus obtained was inserted into plasmid pKM005 at its XbaI site in the possible two orientations. Plasmids III and V, an approximately 300 bp XbaI-XbaI fragment was isolated from plasmid pCX28 (Fig. 1a) and inserted into plasmid pKM005 at its XbaI site in the two possible orientations. These plasmids (I-V) were transformed into a *lacZ* deletion strain SB4288 (F<sup>-</sup> recA thi-I relA mal24 spc12 supE-50 proB lac), and those  $\beta$ -galactosidase activities were tested on MacConkey plates (Difco). Results are shown as LacZ<sup>+</sup> or LacZ<sup>-</sup>. Ability of these clones to inhibit the expression of OmpF protein are also shown as MicF<sup>+</sup> or MicF<sup>-</sup>.

The resulting plasmid, pMYI50 (Fig. 1b) contains the entire coding region of the *ompC* gene and approximately 500 bp of upstream sequences including the *ompC* promoter and the DNA encoding the 5'-end untranslated region of *ompC* mRNA. In order to obtain an *ompC* promoter fragment of different sizes, pMYI50 was digested by Bal31 nuclease at the unique BglII site, followed by the addition of XbaI linkers. The plasmids constructed in this manner carry XbaI fragments that vary in size due to the position of the XbaI site furthest from the Sall site (see Fig. 1b). The different XbaI fragments were subsequently transferred to a promoter-cloning vector, pKM005 which can express the *lacZ* gene only when a promoter fragment is inserted in the right orientation into its unique XbaI site. These experiments revealed that transcription of the *ompC* gene initiates at a site located between 390 and 440 bp downstream from the upstream XbaI site (originally HpaI site). Surprisingly, E. coli transformed with these pKM005 derivatives, including the clone of the shortest XbaI fragment of only 300 bp, CX28 (subcloned from pCX28; Fig. 1a and

b, lost the ability to produce OmpF protein. OmpF protein was clearly produced in the host cells (ompB<sup>+</sup> ompF<sup>+</sup> ompC<sup>+</sup>), while the same cells carrying the clone of the CX28 fragment were not able to produce OmpF protein. The same effect could be observed with cells harboring a clone of a longer fragment such as plasmid I in Fig. 1c. In this clone the lacZ gene was fused immediately after the initiation codon of the ompC gene resulting in the LacZ<sup>+</sup> phenotype of the cells carrying this plasmid. However, when the XbaI-MspI fragment of 87 bp was removed from plasmid I, the cells carrying the resulting plasmid (plasmid II in Fig. 1c) were able to produce OmpF protein. It should be mentioned that a similar DNA fragment of 430 bp in length containing the upstream region of the ompF gene did not block the production of both OmpF and OmpC proteins.

## DNA Sequence Homology Between CX28 and the ompF Gene

The results described above demonstrate that the stretch of DNA approximately 300 bp long, located upstream of the ompC promoter, is able to block ompF expression. In order to elucidate the function of this DNA fragment (CX28), the DNA sequence of this region was determined.

Reference is now made to Fig. 2 which shows the nucleotide sequence of the promoter region and upstream of the ompC gene. Restriction DNA fragments prepared from pMYIII or pMYI50 were labeled at their 3'-end by the method of Sakano et al., *Nature*, 280, 288-294 (1979), using [ $\alpha$ -<sup>32</sup>P] dNTP's and DNA polymerase I large fragment (Klenow fragment). Singly end-labeled DNA fragment was obtained by digestion with a second restriction enzyme. DNA sequence were determined by the method of Maxam and Gilbert, *Methods in Enzymology* 65, 499-560 (1981), using 20%, 10% and 6% polyacrylamide gels in 7 M urea. The RNA polymerase recognition site (-35 region) and the Pribnow box (-10 region) for the ompC and micF promoter, as well as the initiation codon of the ompC gene are boxed. The transcriptional initiation sites determined by SI nuclease mapping for the ompC and micF genes.

Fig. 2 shows the DNA sequence of 500 bp from the XbaI site (originally HpaI) to the initiation codon, ATG, of the ompC gene. The DNA sequence downstream of residue 88 was determined previously. It was found that the sequence from residue 99 to 180 (Fig. 2) has 70% homology with the 5'-end region of the ompF mRNA which includes the Shine-Dalgarno sequence, the initiation codon, and the codons for the first nine amino acid residues of pro-OmpF protein (bases marked by + are homologous to the ompF sequence). A plausible model to explain the above result is that the 300-bp CX28 fragment (Fig. 1c) contains a transcription unit which is directed towards the region upstream of the ompC gene so that the RNA transcript from this region has a sequence complementary to the ompF mRNA. The hybridization between the two RNAs thus blocks the production of OmpF protein.

### Existence of a New Transcription Unit

To determine whether the CX 28 fragment contained an independent transcription unit oriented in a direction opposite from the ompC gene, the lacZ gene was fused at two different sites within the CX28 fragment. In plasmid V, the CX28 fragment was inserted in the opposite orientation with respect to plasmid III (Fig. 1c). This clone was still fully active in suppressing the production of OmpF protein, although it did not produce  $\beta$ -galactosidase (LacZ<sup>-</sup>) (see Fig. 1c). When the fusion junction was shifted to the MspI site at nucleotide 88 (Fig. 2, also see Fig. 1c), the newly constructed clone (plasmid IV) was capable of producing  $\beta$ -galactosidase. However, this plasmid was no longer able to suppress the production of OmpF protein. Although this plasmid contains additional DNA (approximately 200 bp) at the upstream of the lacZ and the CX28 sequences (from residue 300 to 500; Fig. 2), this should not affect the functions of the CX28 fragment since plasmid V is fully active in the suppression of OmpF protein production. These results demonstrate that there is a transcription unit in the CX28 fragment which is independent from the ompC gene promoter and that the CX28 fragment and the ompC gene are transcribed in divergent directions. The fact that plasmid IV can produce  $\beta$ -galactosidase and plasmid IV does not, indicates that the CX28 transcription unit terminates between residue 1 and 88 (Fig. 1c). In fact, a very stable stem-and-loop structure can form between nucleotides 70 and 92 (arrows with letter a in Fig. 2) which is followed by oligo-[T]. This structure is characteristic of  $\rho$ -factor independent transcription termination sites in prokaryotes. The  $\Delta G$  value for this structure was calculated to be -12.5 Kcal according to Salsner, W., *Cold Spring Harbor Symp. Quant. Biol.* 13, 985-1002 (1977).

The initiation site for the CX28 transcript was positioned at nucleotide 237 (Fig. 2) by SI-nuclease

mapping. This result indicates that the CX28 DNA fragment is transcribed to produce a transcript of 174 nucleotides. This was further proven by Northern blot hybridization. In the RNA preparation extracted from cells carrying plasmid III (Fig. 1c), an RNA species is clearly observed to hybridize with the CX28 fragment, which migrates a little slower than 5S RNA. In the control cells, only a small amount of the same RNA was detected. The size of the RNA (CX28 RNA) was estimated on gel to be approximately 6S which is in very good agreement with the size estimated from the sequence (174 bases).

#### Function of the CX28 RNA

As pointed out earlier, the CX28 DNA fragment has extensive homologies with a portion of the *ompF* gene. Thus, part of CX28 RNA is complementary to the *ompF* mRNA and can form an extremely stable hybrid with the *ompF* mRNA as shown in Fig. 3. The  $\Delta G$  value for this hybrid formation was calculated to be -55.5 Kcal. Forty-four bases of the 5'-end untranslated region of *ompF* mRNA, including the Shine-Delgarno sequence for ribosome-binding and 28 bases from the coding region, are involved in the hybrid formation. This hybrid structure is sandwiched by the two stable stem-and-loop structures of the CX28 RNA; one for the 3'-end  $\beta$ -independent transcription termination signal (loop a) and the other at the 5'-end (loop b). The  $\Delta G$  values for loops a and b were calculated to be -12.5 and -4.5 Kcal, respectively.

Referring now to Fig. 3 of the drawings, there is illustrated therein hybrid formation between *micF* and *ompF* mRNA. The sequence of *micF* RNA corresponds to the sequence from residue 237 to 64 in Fig. 2. The *ompF* mRNA sequence was cited from Inokuchi, K. et al., *Nucleic Acids Res.* 10, 6957-6968 (1982). The  $\Delta G$  values for the secondary structures a, b and c were calculated to be -12.5, -4.5 and +2.9 Kcal, respectively.

In Fig. 3 another loop (loop c) is shown. This loop, however, is unlikely to be formed because of its  $\Delta G$  value (+2.9 Kcal). It appears that the formation of the hybrid blocks the translation of *ompF* mRNA. This would explain why clones carrying the CX28 DNA fragment suppress the production of *OmpF* protein. Thus, CX28 RNA is designated as the mRNA-interfering complementary RNA for *ompF* (*micRNA* for *ompF*) and the gene is designated *micF*. It should be noted that when loop a was eliminated by fusing the *micF* gene with the *lacZ* gene, the *MicF* function was abolished (plasmid IV, Fig. 1c). This may be due to the stability of the *micF* RNA or alternatively due to the requirement of loop a for the *MicF* function.

It seemed of interest to examine whether the *micF* gene is under the control of the *ompB* locus as is the *ompC* gene. Various *lacZ* clones were therefore put into four different *ompB* mutants. Reference is now made to Table I.

TABLE I

$\beta$ -Galactosidase Activities of Various Promoter-lacZ Gene Fusion Clones in ompB Mutant Strains  
 $\beta$ -Galactosidase Activity (U)

Strains	Plasmids	pKM004 ( <u>lppP-lacZ</u> )	Plasmid I ( <u>ompCP-lacZ</u> )	Plasmid IV ( <u>micP-lacZ</u> )	pOmpFP-A1 ( <u>ompFP-lacZ</u> )
MC4100 (wild type)					
OmpC <sup>+</sup> OmpF <sup>+</sup>		1360	1808	796	2071
MH1160 ( <u>ompR1</u> )					
OmpC <sup>-</sup> OmpF		1415	102	133	43
MH760 ( <u>ompR2</u> )					
OmpC <sup>-</sup> OmpF <sup>+</sup>		1219	21	102	1521
MH1461 ( <u>envZ</u> )					
OmpC <sup>+</sup> OmpF <sup>-</sup>		905	1500	616	1063

Various ompB mutant strains, MC4100 (F-lacVI69 araDI39 rspL thiA tibB relA; wild type), MH1160 [ompBI01 (ampR1) mutant from MC4100] MH760 [ampB427 (ompR2) mutant from MC4100], MH1461 [tpolI (envZ) mutant from MC4100] were transformed by various promoter-lacZ gene fusion clones. Cells were grown in 10 ml of nutrient broth at 37°C to Klett unit of 1.2. 100  $\mu$ l of the cultures were used for  $\beta$ -galactosidase activity measurement according to the method of Miller, H.J., in Experiments of Molecular Genetics (ed. Miller, H.J.) 352-355 (Cold Spring Harbor Laboratory, New York (1972)). Plasmid pKM004 was derived from pKM005 and pKM004 contains the lpp (the gene for outer membrane lipoprotein) promoter fused to the lacZ gene. Plasmid I and IV are described in Fig. 1c. Plasmid pOmpFP-A1 contains the lacZ gene under the control of the ompF promoter.

As shown in Table I, the lacZ gene under micF control (plasmid IV in Fig. 1C) produces  $\beta$ -galactosidase



in the same manner as the lacZ gene under ompC promoter control (plasmid I in Fig. 1C): high  $\beta$ -galactosidase activity was found in both the wild type and envZ<sup>-</sup> strains but low activity was observed in ompR<sup>-</sup> and ompR2<sup>-</sup> mutants. On the other hand, the lacZ gene under the control of the ompF promoter was not expressed in the ompR<sup>-</sup> cells. In addition, lacZ gene under the control of the lipoprotein promoter, used as a control, was expressed in all strains. These results indicate that the micF gene is regulated by the ompB locus in the same fashion as the ompC gene. It is interesting to note that the lacZ gene under the control of the ompF promoter is constitutively expressed in the envZ<sup>-</sup> (OmpC<sup>+</sup> OmpF<sup>-</sup>) strain. This suggests that the OmpF<sup>-</sup> phenotype of this envZ<sup>-</sup> strain is due to the inhibition of translation of the ompF mRNA by micRNA.

## Promoters of the micF and ompC Genes

Since both the micF and ompC genes appear to be regulated by the ompB locus, the promoters of these genes should have sequence homologies. In order to search for the homologies, the transcription initiation site for the ompC gene was first determined by SI-nuclease mapping. Major transcription initiation takes place at the T residues at position 410 and 411 (Fig. 2; also see Fig. 4).

In Fig. 4 there is shown the homologous sequences between the micF and the ompC genes. Nucleotide numbers correspond to those in Fig. 2. The sequences in a box show the homologous sequences between the two genes. Bars between the two sequences indicate the identical bases. The arrows indicate the transcription initiation sites. The -10 and -35 regions are underlined.

Thus, -10 regions for the micF and ompC genes are assigned as AATAAT (nucleotides 250 to 245 in Fig. 2) and GAGAAT (nucleotides 400 to 405 in Fig. 2), respectively (Fig. 4), both of which show good homology to the consensus sequence, TATAAT. RNA polymerase recognition sites, (-35 regions), for the micF and ompC genes are also assigned as TAAGCA and TTGGAT, respectively (Fig. 4), both of which show 50% homology to the consensus sequence, TTGACA. However, no significant sequence homologies are found between the micF promoter of 63 bp (nucleotides 300 to 238) and the ompC promoter (nucleotides 301 to 409 in Fig. 2). On the other hand, homologous sequences are found in the 5'-end regions of both the transcripts as shown in Fig. 4. Twenty-eight out of 44 bases are homologous (64% homology), and these regions are probably the sites recognized by OmpR protein. It is interesting to note that a homologous sequence to these sequences has also been found in the 5'-end untranslated region of ompF mRNA. Binding experiments of the micF gene and the ompC gene with purified OmpR protein are now in progress.

As indicated hereinabove, regulation of gene expression in *E. coli* is generally controlled at the level of transcription. It has been well established that expression of some genes are suppressed by their specific repressors or activated by their specific inducers. Positive protein factors such as cAMP receptor protein and OmpR protein are also known to regulate gene expression at the level of transcription. Another transcriptional regulatory mechanism is attenuation which plays an important role in controlling expression of operations involved in the biosynthesis of various amino acids of other compounds, see Kolter, R. & Yanofsky, C. *Ann. Rev. Genet.* 16, 113-134 (1982).

In addition, some proteins have been shown to regulate gene expression at the level of translation. The results herein demonstrate the regulation of bacterial gene expression at the level of translation by means of a complementary RNA factor to the translational start region. This novel regulatory mechanism mediated by micRNA is illustrated in Fig. 5.

Fig. 5 illustrates a possible model for the role of micF RNA. OmpR protein binds to the ompF gene under the low osmolarity and promotes the production of OmpF protein. Under the high osmolarity, OmpR protein binds to both the micF and the ompC genes. The micF RNA thus produced hybridizes with the ompF mRNA to arrest its translation.

The possibility that micRNA blocks the expression of the ompF gene at the level of transcription has not been ruled out. However, this is highly unlikely since the lacZ gene fused with the ompF promoter was expressed in the envZ<sup>-</sup> cells (OmpC<sup>+</sup> OmpF<sup>-</sup>; Table 1). In this case lacZ expression is probably due to the inability of lacZ mRNA transcribed from the clone to form a stable hybrid with micRNA. Furthermore, if micRNA is able to bind the nonsense strand of the ompF gene, it would more likely stimulate gene expression rather than block it, since the binding would make the ompF gene more accessible to RNA polymerase.

Regulation by micRNA appears to be an extremely efficient way to block production of a specific

protein without hampering other protein production. At present, the relative ratio between micRNA and ompC production is not known ( $\beta$ -galactosidase activities in Table I do not necessarily reflect their accurate promoter activities, since the promoter regions were not inserted in the same fashion, see Fig. 1c). However, it is reasonable to assume that the micRNA and the ompC are produced coordinately. Therefore, when OmpC protein is produced, micRNA is produced in the same manner. micRNA then blocks the production of OmpF protein proportionally, so that the total amount of OmpC plus OmpF protein is constant.

The binding of micRNA to the ribosome-binding site and the initiation codon is a very effective way to block the translation of the particular mRNA. A similar mechanism has been proposed to explain a translational block in a mutant of bacteriophage T7. It was suggested that the sequence of the 3'-end of a mutant mRNA hybridizes with its own ribosome-binding site to block translation, see Saito, H. & Richardson, C.C. *Cell* 27, 533-542 (1981). It seems reasonable that the micRNA regulatory system may be a general regulatory phenomenon in *E. coli* and in other organisms including eukaryotes. It is a particularly attractive mechanism to very rapidly stop the formation of a protein or to control the ratio of one protein with another. RNA species may have additional roles in the regulation of various cellular activities. In fact, small RNA species have been shown to be involved in the regulation of DNA replication of some plasmids.

In view of the accompanying disclosure it is seen that there is presented in accordance with the practices of this invention a powerful tool and technique for regulating gene expression. Gene expression in accordance with the practices of this invention is regulated by incorporating in or associating with the genetic material of an organism or cellular material which may possess only its native genetic material or which may have been genetically altered by the deletion of native genetic material or the addition of foreign genetic material, DNA which upon transcription along with the genetic material of said organism or cellular material produces an oligoribonucleotide or polyribonucleotide RNA complementary to and/or capable of hybridizing with an mRNA produced by the genetic material of the organism or cellular material so that expression or translation of said RNA is inhibited or prevented.

The regulation of the gene expression of an organism or cellular material in accordance with the practices of this invention is carried out in a transformed organism or cellular material wherein along with the genetic material of said organism or cellular material there is incorporated therein or associated therewith DNA which upon transcription along with the genetic material of said organism or cellular material produces an oligoribonucleotide or polyribonucleotide RNA complementary to and/or capable of binding or hybridizing with an mRNA produced by the genetic material of said organism or cellular material so that the expression or translation of said mRNA is inhibited or prevented.

In the practices of this invention the DNA material or molecule which upon transcription in a transformed organism or cellular material containing said DNA material or molecule produces an oligoribonucleotide or polyribonucleotide RNA complementary to and/or capable of binding or hybridizing with a mRNA produced by the genetic material of said organism or cellular material, may be incorporated or associated with the genetic material of the organism to be transformed by transforming the organism or cellular material with the DNA material or molecule per se directly or by incorporating the DNA material in a plasmid or virus or viral vector and then transforming the organism or cellular material with the plasmid and/or viral vector. The DNA material or molecule may be inserted directly into the nucleus containing the genetic material of the organism or cellular material. The DNA material or molecule effecting transformation of the organism or cellular material may be inserted into the organism through the membrane thereof into the cytoplasm or fluid content of the organism or cellular material into association with the genetic or chromosomal DNA material characterizing the organism. Where desired, convenient or practical, microinjection may be employed to insert the DNA material or molecule into the organism or cellular material to be transformed, such as into the nucleus or cytoplasm of the organism. It is usually convenient to incorporate or associate the DNA material or molecule with the genetic material of the organism or cellular material to be transformed by transfer of the DNA material or molecule through the membrane encompassing the organism or cellular material.

#### Construction of an Artificial Mic Gene

The micF gene produces a 174-base RNA that blocks production of the OmpF protein. This small RNA has two stem-and-loop structures, one at the 3'-end and the other at the 5'-end. Since these structures are considered to play an important role for the function of the micRNA, it was attempted to use these features in the construction of an artificial mic system using the gene for the major outer membrane lipoprotein (lpp) cloned in an inducible expression vector, pIN-II, see Nakamura et al, "Construction of Versatile Expression Cloning Vehicles Using the Lipoprotein Gene of *Escherichia coli*", *EMBO J.* 1, 771-775 (1982). pIN-II vectors are high expression vectors that have the  $\text{lac}^{\text{P}}$  downstream of the lipoprotein promoter, thus allowing high

level inducible expression of an inserted gene. The pIN-II promoter was fused to the *lpp* gene at a unique XbaI site immediately upstream of the Shine-Dalgarno sequence of the *lpp* mRNA. The resulting plasmid was designated as pYMI40. When the expression of the *lpp* gene, in pYMI40, is induced by isopropyl- $\beta$ -D-thiogalactoside (IPTG), a lac inducer, the RNA transcript derived from the *lpp* gene has a possible stem-and-loop structure (at the 5' end). Immediately upstream of the unique XbaI site, see Fig. 6-A, is another stable stem-and-loop structure at its 3' end. The latter loop is derived from the p-independent transcription termination signal of the *lpp* gene. The construction of a general mic cloning vector, pJDC402 was achieved by removing the DNA fragment in pMHO44 between the two loops as shown in Fig. 6 -A. An RsaI site immediately upstream of the termination site was changed to an EcoRI site by partial digestion of pYMI40 followed by insertion of an EcoRI linker. The resulting plasmid, pMHO44 was partially digested with EcoRI, followed by a complete digestion with XbaI. The single stranded portions of the linear DNA fragment were filled in with DNA polymerase I (large fragment), and then treated with T4 DNA ligase, resulting in the formation of the plasmid, pJDC402, which lost the fragment between the XbaI and the RsaI sites. As a result of this procedure, both an EcoRI and an XbaI site were recreated at the junction. Thus the unique XbaI site can serve as the insertion site for any DNA fragment, and the RNA transcript from the artificial mic gene produces an RNA which has a similar structure to the *micF* RNA; the portion derived from the inserted DNA is sandwiched by two loop structures, one at the 5' and one at the 3'-end.

The following is a more detailed description of Fig. 6-A and Fig. 6-B. As illustrated in Fig. 6-A for the construction of pJDC402, restriction sites are indicated as follows: X, XbaI; P, PvuII; E, EcoRI. *lpp*<sup>P</sup> and *lac*<sup>PO</sup> are the lipoprotein promoter and the lactose promoter operator, respectively. Amp<sup>r</sup> is the Ampicillin resistance gene. Cross hatches represent the lipoprotein promoter. Solid dots represent the lactose promoter operator. Slashes indicate the lipoprotein signal sequence, and the solid bar represents the coding region for the mature portion of the lipoprotein. The open dots represent the transcription termination region derived from the *lpp* gene. The open bar represents the 5' nontranslated region of the lipoprotein mRNA.

In Fig. 6-B for the construction of *mic(lpp)* pJDC4I2, open arrows represent promoters. The PvuII site was converted to an XbaI site by inserting an XbaI linker (TCTAGAG). This fragment was inserted into the unique XbaI site of pJDC402 in the reverse orientation forming pJDC4I2. a and b show the *mic(lpp)* RNAs initiating at the *lpp* and *lac* promoters, respectively.

### 30 Construction of the *mic(lpp)* Gene

Using this mic cloning vector, pJDC402, it was first attempted to create a mic system for the *lpp* gene of *E. coli*, in order to block the synthesis of the lipoprotein upon induction of the *mic(lpp)* gene. For this purpose it is necessary to first isolate the DNA fragment containing the Shine-Dalgarno sequence for ribosome binding, and the coding region for the first few amino acid residues of prolipoprotein. To do this the PvuII site immediately after the coding region of prolipoprotein signal peptide was changed to an XbaI site by inserting an XbaI linker at this position. The resulting plasmid was then digested with XbaI, and the 112-bp XbaI-XbaI (originally PvuII-XbaI) fragment was purified. This fragment encompassing the Shine-Dalgarno sequence and the coding region for the first 29 amino residues from the amino terminus of prolipoprotein was purified. This fragment was then inserted into the unique XbaI site of pJDC402 in the opposite orientation from the normal *lpp* gene. The resulting plasmid, designated as pJDC4I2, is able to produce *mic(lpp)* RNA, an RNA transcript complementary to the *lpp* mRNA, upon induction with IPTG.

It should be pointed out that another important feature of the mic expression vector, pJDC402, is that it contains a HinfI site immediately upstream of the *lpp* promoter and another one immediately downstream of the transcription termination site. These two HinfI sites can be used to remove a DNA fragment containing the entire mic transcription unit which can then be inserted back into the unique PvuII site of the vector. In this manner, the entire *mic* gene can be duplicated in a single plasmid. One would expect a plasmid containing two identical *mic* genes to produce twice as much micRNA as a plasmid containing a single *mic* gene. Such a plasmid was constructed containing two *mic(lpp)* genes and designated as pJDC422.

### 50 Expression of the *mic(lpp)* Gene

In order to examine the effect of the artificial *mic(lpp)* RNA, cells were pulse-labeled for one minute, with [<sup>35</sup>S]-methionine, one hour after induction of the *mic(lpp)* RNA with 2mM IPTG. The cells harboring the vector, pJDC402, produce the same amount of lipoprotein either in the absence or the presence of the inducer, IPTG, as quantitated by densitometric scanning of the autoradiogram and normalizing. Lipoprotein production was reduced approximately two-fold in the case of cells carrying pJDC4I2 in the absence of IPTG and approximately 16-fold in the presence of IPTG. The reduction in lipoprotein synthesis in the

absence of IPTG is considered to be due to incomplete repression of the mic(lpp) gene. In the case of cells carrying pJDC422, where the mic(lpp) gene was duplicated, lipoprotein production is now reduced 4-fold in the absence of IPTG, and 31-fold in the presence of IPTG. These results clearly demonstrate that the production of the artificial mic(lpp) RNA inhibits lipoprotein production, and that the inhibition is proportional to the amount of the mic(lpp) RNA produced. It should be noticed that the mic(lpp) RNA is specifically blocking the production of lipoprotein, and that it does not block the production of any other proteins except for OmpC protein. The fact that the induction of the mic(lpp) gene reduces the production of the OmpC plus OmpF proteins was found to be due to unusual homology between the lpp and the ompC gene as discussed hereinafter.

There are several mechanisms by which the mic inhibition may occur. One mechanism is that the micRNA binds to the mRNA preventing the ribosome from binding the mRNA. Other possible mechanisms include: destabilization of the mRNA, attenuation of the mRNA due to premature termination of transcription, or inhibition of transcription initiation. If the inhibitory effect of the micRNA is solely at the level of attenuation or transcription initiation one would expect the mic effect to be somewhat delayed due to the fact that the functional half-life of the lipoprotein mRNA is 12 minutes. Therefore, it was examined how rapidly lipoprotein production is inhibited upon induction of the mic(lpp) RNA by pulse-labeling *E. coli* JA221/F<sup>+</sup>lacI<sup>q</sup> harboring pJDC412, with [<sup>35</sup>S]-methionine at various time points after induction with IPTG. It was determined that lipoprotein production was maximally inhibited by 16-fold within 5 minutes after the addition of IPTG. This result indicates that inhibition of lipoprotein production is primarily due to the binding of the mic(lpp) RNA to the lpp mRNA, resulting in the inhibition of translation of the lpp mRNA and/or destabilization of the mRNA.

#### lpp mRNA Production in the Presence of mic(lpp) RNA

It appeared interesting to examine whether the mic(lpp) RNA also affects the level of the lpp mRNA, since the expression of the micF gene substantially reduced the amount of the ompF mRNA. For this purpose, there was isolated total cellular RNA one hour after the induction of the mic(lpp) gene with IPTG. The RNA preparation was analyzed after electrophoresis in a formaldehyde agarose gel and subsequently transferred onto nitrocellulose paper. The paper was then hybridized with a probe specific to the mic(lpp) RNA, or to the lpp mRNA. There was also used a probe specific for the ompA mRNA as an internal control. Again pJDC402 shows no difference in the production of the lpp mRNA in the absence or presence of IPTG. Due to the fact that the double stranded primer used to make the probe for these experiments contains a portion of the lac operon, the probes hybridize to any transcript containing the lac promoter such as the mic(lpp) RNA from JDC412 and the short nonsense transcript from pJDC402. Cells harboring pJDC412 contain a reduced amount of the lpp mRNA in the absence of IPTG and a greatly reduced amount of the lpp mRNA in the presence of IPTG. There was shown the production of the mic(lpp) RNA in the absence and the presence of IPTG in cells harboring pJDC412. Therefore, even in the absence of IPTG, a significant amount of the mic(lpp) RNA is produced, which is consistent with the results of the lipoprotein production observed earlier. The fact that the lpp mRNA disappears upon induction of the mic(lpp) RNA indicates that the mechanism of action of the micRNA is not solely at the level of translation. Tests demonstrated there are two mic(lpp) RNAs of different sizes. The sizes of these transcripts were estimated to be 281 and 197 bases, which correspond to transcripts initiating at the lipoprotein promoter (the larger RNA) and initiating at the lac promoter (the smaller RNA).

#### Inhibition of OmpC Production with the mic(ompC) Gene

It was also possible to achieve an almost complete inhibition of OmpC synthesis by artificially constructing mic(ompC) genes. The first construct, pAM320, carrying two mic(ompC) genes gives rise to an RNA molecule complementary to 20 nucleotides of the leader region and 100 nucleotides of the coding region of the ompC mRNA. This was achieved by changing the unique BglII site in the ompC structural gene and the MnlI site, 20 nucleotides upstream of the ATG initiation codon to XbaI sites. The resulting 128-bp XbaI fragment was then inserted into pJDC402 in the opposite orientation from the OmpC gene and a second copy of the mic(ompC) gene was introduced in a manner similar to that described for the pJDC422 construction. The resulting plasmid, pAM320, has the second mic(ompC) gene inserted in the orientation opposite to the first one. Reversing the orientation of the second mic gene did not change the expression or stability of the plasmid. A second construct, pAM321, was designed to extend the complementarity between the micRNA and the ompC mRNA to include a longer leader sequence than in the case of pAM320, 72 nucleotides of the leader region instead of 20. This plasmid was constructed as described for

pAM320, except that the MnlI site changed to an XbaI site was located 72 nucleotides bp upstream of the *ompC* initiation codon.

Commasie Brilliant Blue stained gel patterns of the outer membrane proteins isolated from *E. coli* JA221/F'lacI<sup>q</sup> harboring the mic cloning vector pJDC402, pAM320 and pAM32I were obtained. The effect of the addition of IPTG was clearly seen by the appearance of  $\beta$ -galactosidase. The induction of the mic(*ompC*) RNA from pAM320 caused a substantial decrease (approximately 5-fold) in *OmpC* production, compared to pJDC402. Induction of the longer mic(*ompC*) RNA from pAM32I decreased *OmpC* synthesis more dramatically (approximately 20-fold compared to pJDC402). *OmpC* production could hardly be detected in the cells harboring pAM32I when they were pulse-labeled for one minute after a one-hour induction with IPTG. In the same experiment, *OmpC* synthesis decreased approximately 7-fold when the mic(*ompC*) gene in cells harboring pAM320 was induced with IPTG. Marked decreases in *OmpC* expression were also observed when plasmids containing single copies of the mic(*ompC*) genes here induced. Again, the longer mic(*ompC*) gene had a greater effect. The increased efficiency of mic-mediated inhibition with pAM320 may indicate that the effectiveness of the micRNA function is related to the extent of complementarity to the 5'-end of the mRNA.

It was interesting to note that the synthesis of either of the mic(*ompC*) RNAs described above caused a decrease not only in *OmpC* synthesis but also in lipoprotein synthesis. This inhibitory effect of the mic(*ompC*) RNA on lipoprotein production appears to be due to the unexpected homology between the *lpp* mRNA sequence and the *ompC* mRNA as illustrated in Fig. 7. This feature explains why pAM320 and pAM32I are exerting a mic effect on lipoprotein production. Such an explanation would predict that induction of the mic(*lpp*) RNA from pJDC412 and pJDC422 should decrease the synthesis of the *OmpC* protein, and this was found to be the case.

In Fig. 7, there is illustrated a region of homology between the *lpp* mRNA (top line) and the *ompC* mRNA (bottom line). Bars connect identical bases. Both mic(*ompC*) RNAs have the potential to hybridize across this homologous region. The Shine-Dalgarno Sequences (S.D.) and AUG initiation codons are boxed.

#### Inhibition of *OmpA* Production with mic(*ompA*) RNA

In an effort to determine what components contribute to the effectiveness of a micRNA, several mic genes were constructed from the *ompA* gene. The *ompA* gene was selected for this because the leader and the coding regions of the *ompA* mRNA have been characterized extensively. Five DNA fragments (see I through V of Fig. 8) were individually cloned into the XbaI site of pJDC402 in the orientation promoting the production of mic(*ompA*) RNAs. The resulting mic(*ompA*) plasmids containing fragments I-V were designated as pAM30I, pAM307, pAM3I3, pAM3I4, and pAM3I8, respectively. Each plasmid contains only one copy of the described mic(*ompA*) gene.

In Fig. 8, the top line shows the structure of the *E. coli ompA* gene. The arrow represents the promoter and the open bar represents the region encoding the 5'-leader region of the *ompA* mRNA. The slashed bar and shaded bar represent the portions of the *ompA* gene encoding the signal sequence and the mature *OmpA* protein, respectively. Restriction fragment I (HphI-HpaI) was inserted into the XbaI site of pJDC402, see Fig. 6-A, in the orientation opposite from that depicted here, as outlined in Fig. 6-B for mic(*lpp*), to create the plasmid, pAM30I. The other mic(*ompA*) plasmids were similarly constructed from: fragment II, pAM307; fragment III, pAM3I3; fragment IV, pAM3I4; fragment V, pAM3I8. The positions of the Shine-Dalgarno sequence (SD), ATG initiation codon (ATG), and relevant restriction sites are shown.

*E. coli* JA221/F'lacI<sup>q</sup> containing each of the mic(*ompA*) plasmids was pulse-labeled with [<sup>35</sup>S]-methionine for one minute with and without a one-hour prior preincubation with IPTG. Electrophoretic patterns of the outer membrane proteins isolated from these cultures were obtained. The autoradiographs revealed that each of the five mic(*ompA*) genes is capable of inhibiting *OmpA* synthesis. The mic(*ompA*) genes appear to be less effective than the mic(*lpp*) and mic(*ompC*) genes described earlier, but this problem was circumvented by increasing the mic(*ompA*) gene dosage.

The plasmid pAM30I, encoding an mRNA complementary to a 258 base region of the *ompA* mRNA encompassing the translation initiation site (fragment I in Fig. 8) was found to inhibit *OmpA* synthesis by approximately 45 percent. A similar inhibition was obtained with pAM307, by approximately 51 percent. This plasmid contains fragment II (see Fig. 8) which does not include any DNA sequences corresponding to the *ompA* structural gene. The inhibition by pAM307 was not surprising because the mic(*ompC*) experiments described earlier showed that increased complementarity to the 5'-leader region of the mRNA was more effective in micRNA-mediated inhibition. On the other hand, pAM3I3, which produces a micRNA that is complementary only to the portion of the *ompA* structural gene covered by fragment III (See Fig. 8 which spans the coding region for amino acid residues 4 through 45 of pro-*OmpA*), was also effectively able to

inhibit OmpA synthesis by approximately 54 percent, indicating that the micRNA does not need to hybridize to the initiation site for protein synthesis and/or to the 5'-leader region of the target mRNA in order to function. This was also confirmed using mic(lpp) genes. Two mic(lpp) RNAs which were complementary to only the coding region of the lpp mRNA have also been found to inhibit lipoprotein production. The effect of the mic(lpp) genes in pJDC4I3 and pJDC4I4 which were constructed from the lpp structural gene fragments coding for amino acid residues 3 to 29, and 43 to 63 of prolipoprotein, respectively, were observed. Both pJDC4I3 and pJDC4I4, however, exhibit only a 2-fold inhibition of lipoprotein synthesis indicating that a DNA fragment covering the translation initiation site (which caused a 16-fold inhibition) is more effective in the case of the mic(lpp) genes.

Fragment IV (see Fig. 8) was chosen to test the effectiveness of a micRNA complementary only to the 5' leader region of the ompA mRNA. The resulting construct pAM3I4, synthesizes a micRNA complementary to a 68-base stretch of the ompA mRNA leader located 60 bases upstream of the AUG initiation codon. pAM3I4 exhibits a very weak mic effect, inhibiting OmpA synthesis by only about 18 percent. The significant differences in the mic effect between fragments II and IV (see Fig. 8) clearly demonstrates that the complementary interaction within the region of the mRNA that interacts with the ribosome i.e., the Shine-Dalgarno sequence and/or the coding region, is very important for the effective mic function, although it is not absolutely required. It is also interesting to note that shortening the mic(ompA) gene from fragment I to V had little effect on its efficiency, a 45 percent compared to a 48 percent decrease, respectively.

In order to construct a plasmid capable of inhibiting OmpA synthesis more effectively than those discussed above, plasmids were constructed containing more than one mic(ompA) gene. These plasmids, pAM307 and its derivatives pAM3I9 and pAM3I5 were compared. The latter two plasmids contain two and three copies of the mic(ompA) gene in pAM307, respectively. While pAM307 inhibited OmpA synthesis by approximately 47 percent, pAM3I5 and pAM3I9 inhibited OmpA synthesis by 69 percent and 73 percent, respectively.

The results presented hereinabove clearly demonstrate that the artificial mic system and techniques of this invention can be used for specifically regulating the expression of a gene of interest. In particular, the inducible mic system for a specific gene is a novel and very effective way to study the function of a gene. If the gene is essential, conditional lethality may be achieved upon the induction of the mic system, somewhat similar to a temperature-sensitive mutation. It should be noted, however, that the mic system blocks the synthesis of the specific protein itself while temperature-sensitive mutations block only the function of the protein without blocking its synthesis.

From this invention, the following has become evident:

(a) The production of an RNA transcript (micRNA) that is complementary to a specific mRNA inhibits the expression of that mRNA.

(b) The production of a micRNA specifically blocks the expression of only those genes which share complementarity to the micRNA.

(c) The induction of micRNA production blocks the expression of the specific gene very rapidly in less than the half-life of the mRNA.

(d) The micRNA also reduces the amount of the specific mRNA in the cell, as was found when the natural micF gene was expressed, as well as when the artificially constructed mic(lpp) gene was expressed in the present invention.

(e) There is a clear effect of gene dosage; the more a micRNA is produced, the more effectively the expression of the target gene is blocked.

In the practices of this invention, it appears that micRNAs complementarity to regions of the mRNA known to interact with ribosomes are the most effective. Using the lpp gene as an example, it appears that a mic(lpp) RNA that can hybridize to the Shine-Dalgarno sequence and the translation-initiation site of the lpp mRNA inhibits lipoprotein synthesis more efficiently than one which cannot. However, for the ompA gene, micRNAs complementarity to both the Shine-Dalgarno sequence and the translation-initiation site, just the Shine-Dalgarno sequence, or the structural gene alone were equally effective.

For some genes, such as ompC and lpp, the region of the gene encompassing the translation-initiation site may not contain a unique sequence, and micRNA induction results in the inhibition of the production of more than one protein. In these cases, another region of the gene may be used to construct the mic gene. The length of the micRNA is another variable to be considered. The longer mic(ompC) RNA was 4-fold more effective at inhibiting OmpC production than the shorter mic(ompC) RNA. It should be noted that the inhibition of lipoprotein expression by the mic(ompC) RNA was less effective with the longer mic(ompC) RNA, in spite of the fact that the region of the two mic(ompC) RNAs complementary to the lipoprotein mRNA is the same. This indicates that higher specificity may be achieved by using longer micRNAs. In contrast to the mic(ompC) genes, length did not appear to be a significant factor for the mic(ompA) RNA-

mediated inhibition of OmpA production. In addition, the secondary structure of the micRNA most likely plays an important role in micRNA function.

There are several mechanisms by which the micRNA may function to inhibit expression of the specific gene. It is most likely that the micRNA primarily acts by binding to the mRNA, thereby preventing the interaction with ribosomes as proposed earlier. This hypothesis is supported by the fact that the mic(lpp) RNA inhibited lipoprotein production much faster than the time expected if only transcription was affected based on the half-life of the lpp mRNA. Concerning how micRNA causes a reduction in the amount of lipoprotein mRNA, a plausible model to explain this reduction is that the mRNA is less stable when ribosomes are not traversing the entire mRNA. Another possible model to explain this reduction in mRNA level is that complementary hybrid formation between the micRNA and the mRNA causes premature termination of transcription or destabilization of the mRNA. Alternatively, the micRNA may directly inhibit the initiation of transcription, or cause pausing of mRNA elongation in a manner similar to that described for the function of a small complementary RNA species in ColEI replication, see Tomizawa et al., The importance of RNA secondary structure in ColEI primer formation. Cell 31, 575-583 (1982).

The mic system of this invention has great potential in its application, in prokaryotic as well as eukaryotic cells, to block, permanently or upon induction, the expression of various toxic or harmful genes such as drug resistance genes, oncogenes, and phage or virus genes and the expression of other genes.

In the development and demonstration of the practices of this invention as described herein, the following materials and procedures were employed.

#### Strain and Medium

E. coli JA221 (hsdr leuB6 lacY thi recA ΔtrpE5)F'(lacI<sup>q</sup> proAB lacZYA) was used in all experiments. This strain was grown in M9 medium (J.H. Miller, Experiments in Molecular Genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (1972)) supplemented with 0.4 percent glucose, 2 μg/ml thiamine, 40 μg/ml each of leucine and tryptophan, and 50 μg/ml ampicillin, unless otherwise indicated.

#### Materials

Restriction enzymes were purchased from either Bethesda Research Laboratories or New England BioLabs. T4 DNA ligase and E. coli DNA polymerase I (large fragment) were purchased from Bethesda Research Laboratories. All enzymes were used in accordance with the instructions provided by the manufacturer. XbaI linkers (CTCTAGAG) were purchased from New England BioLabs.

#### DNA Manipulation

Plasmids pJDC402, pJDC412, and pJDC422 were constructed as described herein and Fig. . Plasmids pJDC413 and pJDC414 were constructed by isolating the 80-bp AluI fragment from the lpp gene encoding amino acid residues 3 through 29 of prolipoprotein for pJDC413 and the 58-bp AluI fragment encoding amino acid residues 43 through 63 of prolipoprotein for pJDC414. The fragments were blunt end ligated into pJDC402 which was first digested with XbaI followed by treatment with DNA polymerase I (large fragment).

The isolation of the appropriate ompC fragments for mic(ompC) construction involved a subcloning step due to the absence of suitable unique restriction sites between the ompC promoter and structural gene. Two derivatives of the ompC containing plasmid, pMY150, lacking either the 471-bp XbaI-MnII ompC promoter containing fragment (pDR001 and pDR002, respectively), but containing an XbaI site in its place, were isolated. The unique BglII sites in each of these plasmids were changed to XbaI sites by treatment with DNA polymerase I (large fragment) and ligation with synthetic XbaI linkers. Following XbaI digestion, the 123-bp XbaI fragment from pDR001 and the 175-bp XbaI fragment from pDR002 were individually isolated and cloned into the XbaI site of pJDC402 to create pAM308 and pAM309, respectively. pAM320 contains the HindIII fragment covering the mic(ompC) gene isolated from pAM308 cloned into the PvuII site of pAM308. pAM321 was similarly constructed from pAM309 to also contain two mic(ompC) genes.

The mic(ompA) plasmids pAM301, pAM307, pAM313, pAM314, and pAM318 were constructed as described in a manner similar to the construction of the mic(lpp) and the mic(ompC) genes. To construct pAM319, the HindIII fragment containing the mic(ompA) gene was isolated from pAM307 and inserted back into the PvuII site of pAM307. pAM315 was constructed in the same manner as pAM319 except that it contains two HindIII fragments inserted into the PvuII site of pAM307.

#### Analysis of outer membrane protein production

E. coli JA221/F<sup>+</sup>lacI<sup>q</sup> carrying the appropriate plasmid were grown to a Klett-Summerson colorimeter reading of 30, at which time IPTG was added to a final concentration of 2 mM. After one additional hour of growth (approximately one doubling), 50  $\mu$ Ci of [<sup>35</sup>S]-Methionine (Amersham, 1000 Ci/mMole) was added to 1 ml of the culture. The mixture was then incubated with shaking for one minute, at which time the labeling was terminated by addition of 1 ml ice cold stop solution (20 mM sodium phosphate, [pH 7.1] containing 1 percent formaldehyde, and 1 mg/ml methionine). Cells were washed once with 10 mM sodium phosphate, pH 7.1, suspended in 1 ml of the same buffer, and sonicated with a Heat Systems Ultrasonics sonicator model W-220E with a cup horn adapter for 3 minutes (in 30 second pulses). Unbroken cells were removed by low speed centrifugation prior to collecting the outer membrane. Cytoplasmic membranes were solubilized during a 30 minute incubation at room temperature in the presence of 0.5 percent sodium lauroyl sarcosinate and the outer membrane fraction was precipitated by centrifugation at 105,000 X g for 2 hours.

Lipoprotein and OmpA were analyzed by Tris-SDS polyacrylamide gel electrophoresis (SOS-PAGE). To analyze OmpC production, urea-SDS polyacrylamide gel electrophoresis (urea-SDS-PAGE) was used. Proteins were dissolved in the sample buffer and the solution was incubated in a boiling water bath for 8 minutes prior to gel application. The autoradiographs of dried gels were directly scanned by a Shimadzu densitometer. To determine relative amounts of the band of interest, the ratio of the area of the peak of interest to the area of an unaffected protein peak, was determined for each sample.

#### RNA Analysis

Cells were grown and labeled with [<sup>3</sup>H]-uridine, then cell growth was stopped by rapidly chilling the culture on ice for less than 5 minutes. The cells were collected by centrifugation at 8000 rpm for 5 minutes. RNA was isolated using the following procedure. The cells were quickly resuspended in hot lysis solution (10 mM Tris-HCl [pH 8.0], 1 mM EDTA, 350 mM NaCl, 2 percent SDS and 7 M urea) with vigorous vortexing for 1 minute. The mixture was immediately extracted, twice with phenol:chloroform (1:1) and twice with chloroform alone. One tenth volume of 3 M sodium acetate (pH 5.2) was added to the mixture and 3 volumes of ethanol was added to precipitate the RNA. The precipitate was then dissolved in TE buffer (10 mM Tris-HCl [pH 7.5], 1 mM EDTA). For gel electrophoresis, equal counts were loaded in each lane. The RNA was separated on a 1.5 percent agarose gel containing 6 percent formaldehyde. The running buffer was 20 mM MOPS (3-[N-morpholino]propanesulfonic acid [Sigma]), 5 mM sodium acetate and 1 mM EDTA, pH 7.0.

RNA was transferred to nitrocellulose paper. MI3 hybridization probes specific for the mic(lpp) RNA and lpp mRNA were individually constructed by cloning the 112-bp XbaI fragment shown in Fig. 1-B into MI3 mp9 in the appropriate orientation. A probe specific for the ompA mRNA was constructed by inserting a 1245-bp XbaI-EcoRI fragment (originally an EcoRV-PSTI fragment) into MI3 mpl0 and the probes were labeled.

#### Claims

1. A non-human organism containing a nucleic acid construct, said nucleic acid construct being an artificial nucleic acid construct which, upon introduction into a cell of said organism containing a gene, regulates the function of said gene and produces a ribonucleotide sequence which does not naturally occur in the cell, is complementary to at least a portion of a ribonucleotide sequence transcribed by said gene, and regulates the function of said gene.
2. An organism containing a nucleic acid construct, said nucleic acid construct being an artificial nucleic acid construct which, upon introduction into a cell of said organism containing a gene, regulates the function of said gene and produces a ribonucleotide sequence which does not naturally occur in the cell, is complementary to at least a portion of a ribonucleotide sequence transcribed by said gene, and regulates the function of said gene, provided that said organism is not exempt under Articles 53a) and 53b) EPC.
3. A non-human organism containing a nucleic acid construct, said nucleic acid construct being an artificial nucleic acid construct which, upon introduction into a cell of said organism containing a gene, regulates the function of said gene, and which contains the following nucleic acid segments:
  - (a) a transcriptional promotor segment;
  - (b) a transcription termination segment; and
  - (c) a segment of said gene, said gene segment located between said promotor segment and said termination segment and being inverted with respect to said promoter segment and said termination



segment, the polarity of said inverted gene segment being the same as that of said promoter segment and said termination segment, whereby transcription of the inverted gene segment occurs in a direction opposite to the direction of transcription of the gene to thereby regulate the function of said gene.

5

4. An organism containing a nucleic acid construct, said nucleic acid construct being an artificial nucleic acid construct which, upon introduction into a cell of said organism containing a gene, regulates the function of said gene, and which contains the following nucleic acid segments:

10

- (a) a transcriptional promoter segment;
- (b) a transcription termination segment; and
- (c) a segment of said gene, said gene segment located between said promoter segment and said termination segment and being inverted with respect to said promoter segment and said termination segment, the polarity of said inverted gene segment being the same as that of said promoter segment and said termination segment, whereby transcription of the inverted gene segment occurs in a direction opposite to the direction of transcription of the gene to thereby regulate the function of said gene, provided that said organism is not exempt under Article 53a) and 53b) EPC.

15

5. The organism of any one of claims 1 to 4, wherein said nucleic acid construct or nucleic acid sequence segment encodes a ribonucleotide sequence complementary to a 5' end non-coding portion of said ribonucleotide acid sequence transcribed by said gene.

20

6. The organism of any one of claims 1 to 4, wherein said nucleic acid construct or nucleic acid sequence segment encodes a ribonucleotide sequence complementary to a ribosome binding portion of said ribonucleotide acid sequence transcribed by said gene.

25

7. The organism of any one of claims 1 to 4, wherein said nucleic acid construct or nucleic acid sequence segment encodes a ribonucleotide sequence complementary to the translation initiation portion of said ribonucleotide acid sequence transcribed by said gene.

30

8. The organism of any one of claims 1 to 7, wherein said gene is an oncogene.

9. The organism of any one of claims 1 to 7, wherein said gene is a viral gene.

10. The organism of any one of claims 1 to 9, wherein said gene encodes a protein.

35

11. The organism of any one of claims 1 to 10, wherein said transcriptional promoter segment comprises an inducible promoter.

12. The organism of any one of claims 1 to 11, wherein said gene segment includes the 5' non-coding region of said gene.

40

13. The organism of any one of claims 1 to 12, wherein said gene segment includes the ribosome binding portion of said gene.

45

14. The organism of any one of claims 1 to 13, wherein said gene segment includes the translation initiation portion of said gene.


15. The organism of any one of claims 1 to 14 containing said artificial nucleic acid construct incorporated in or associated with the chromosomal genetic material.

50

16. The organism according to claim 15 containing said artificial nucleic acid construct incorporated in or associated with the chromosomal genetic material in the nucleus of its cellular material.

55

FIG. 1



**CompC**




FIG. 3




FIG. 4




FIG. 5




Figure 6a





Figure 6b





**FIG. 7**





FIG. 8





European  
Patent Office

## EUROPEAN SEARCH REPORT

Application Number

EP 91 11 1983

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X,D	CELL, vol. 34, September 1983, pages 683-691, MIT; R.W. SIMONS et al.: "Translational control of IS10 transposition" * Whole document *	1-16	C 12 N 1/21 // C 12 N 15/63
X,P,D	PROCEEDINGS OF THE JAPAN ACADEMY, series B, vol. 59, December 1983, pages 335-338; T. MIZUNO et al.: "Regulation of gene expression by a small RNA transcript (micRNA) in Escherichia coli K.12" * Whole document *	1-16	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C 12 N
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of search 29 August 91	Examiner CUPIDO M.
<div>CATEGORY OF CITED DOCUMENTS</div> <div>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention</div> <div>E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</div>			